



## Faunal mortality associated with massive beaching and decomposition of pelagic *Sargassum*



R.E. Rodríguez-Martínez<sup>a,\*</sup>, A.E. Medina-Valmaseda<sup>a</sup>, P. Blanchon<sup>a</sup>, L.V. Monroy-Velázquez<sup>b</sup>, A. Almazán-Becerril<sup>c</sup>, B. Delgado-Pech<sup>c</sup>, L. Vásquez-Yeomans<sup>d</sup>, V. Francisco<sup>e</sup>, M.C. García-Rivas<sup>e</sup>

<sup>a</sup> Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, Prolongación Avenida Niños Héroes S/N, Puerto Morelos, Quintana Roo 77580, Mexico

<sup>b</sup> Colección Nacional de Crustáceos, Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad Universitaria, Coyoacán, Ciudad de Mexico 04510, Mexico

<sup>c</sup> Laboratorio Ecología Costera, Unidad de Ciencias del Agua, CICY, Calle 8, No. 39, S. M. 64, Cancún, Quintana Roo, Mexico

<sup>d</sup> El Colegio de la Frontera Sur, Unidad Chetumal, Depto. de Sistemática y Ecología Acuática, Av. Centenario km 5.5, Col. Pacto Obrero, 77014 Chetumal, Quintana Roo, Mexico

<sup>e</sup> Parque Nacional Arrecife de Puerto Morelos, Comisión Nacional de Áreas Naturales Protegidas, Prolongación Avenida Niños Héroes S/N, Puerto Morelos, Quintana Roo 77580, Mexico

### ARTICLE INFO

#### Keywords:

Macroalgae bloom  
Fauna mortality  
Mexican Caribbean

### ABSTRACT

In 2018, the Mexican Caribbean coast received a massive influx of pelagic *Sargassum* spp. that accumulated and decayed on beaches producing organic decomposition products that made the water turbid and brown. Between May and September of the same year there were several reports of mass mortality of marine biota in this area. From these reports we estimate that organisms belonging to 78 faunal species died as result of this event, with demersal neritic fish and Crustacea being the most affected groups. The cause of mortality appears to be the combined effect of high ammonium and hydrogen sulfide concentrations, together with hypoxic conditions. If massive arrival of pelagic *Sargassum* spp. continues and algae is left to decay on the beach in large volumes then deterioration in water quality could affect coral reefs close to shore. Furthermore, barriers placed in lagoons to intercept the *Sargassum* spp. before it reaches the beach could impact reef fauna if the algae is left to die and sink on site.

Mass mortality events of coastal marine fauna have been reported worldwide due to several factors including extreme temperature changes (Hsieh et al., 2008), resuspension of anoxic-hypoxic or toxic sediment (Justić et al., 1996), emerging diseases (Lessios et al., 1984; Harvell et al., 1999) and micro- and macroalgal blooms (Anderson, 2007). Furthermore, over the past few decades, macroalgae blooms have increased worldwide (Lapointe, 1997), posing a major threat to public health, ecosystem health, and to fisheries and economic development (Anderson, 2007). Mass mortality events of fish, associated with brown and green macroalgae blooms, have been reported from several coasts, such as those off Brazil (Pinheiro et al., 2010; Sissini et al., 2017), Nigeria (Oyesiku and Egunyomi, 2014; Adet et al., 2017), and the Ivory Coast (Sankaré et al., 2016).

Here we report a faunal mass-mortality event along the Mexican

Caribbean coast in 2018 associated with the massive influx of pelagic *Sargassum* spp. and its subsequent decay that resulted in decomposition products that made the water brown and turbid. This turbid water extended for hundreds of meters from the shore (Supp. Fig. 1) and was reported as a *Sargassum*-brown-tide by van Tussenbroek et al. (2017) to differentiate it from the terms “golden floating rainforest” (Laffoley et al., 2011) and “golden tides” (Smetacek and Zingone, 2013), which refer to drifting masses of pelagic *Sargassum* spp. in the open ocean. Historically this shoreline has received relatively small quantities of pelagic *Sargassum* spp. that presumably drifted from the Sargasso Sea through the northern passages of the Caribbean. In 2014, pelagic *Sargassum* (*Sargassum fluitans* and *S. natans*) started to arrive to the Mexican Caribbean coast in unusually large quantities, reaching a peak in September 2015, when  $\sim 2360 \text{ m}^3$  algae  $\text{km}^{-1}$  washed ashore between

\* Corresponding author.

E-mail address: [rosaer@cmarl.unam.mx](mailto:rosaer@cmarl.unam.mx) (R.E. Rodríguez-Martínez).




Fig. 1. Relative abundance of five groups of fauna observed dead along the Mexican Caribbean coast during the 2018 massive influx of *Sargassum* spp. event.

Cancun and Puerto Morelos (Rodríguez-Martínez et al., 2016). After a decrease in 2016 and 2017, the influx of *Sargassum* spp. resumed in 2018, peaking in May at  $8793 \text{ m}^3 \text{ km}^{-1}$  (SD:  $12,485 \text{ m}^3 \text{ km}^{-1}$ ) (Unpublished data). Satellite-derived observations of drifting *Sargassum* spp. masses (Gower et al., 2013) and hind cast models of *Sargassum* spp. landfalls (Johnson et al., 2013) suggest an origin off the coast of Brazil. In 2015, the *Sargassum*-brown-tide extended for over 200 m from shore at some sites (Supp. Fig. 1) and caused mortality of seagrasses and corals (van Tussenbroek et al., 2017). During this event, however, dead fauna were rarely observed along the beach, and no reports of mass mortality were made.

In 2018, between May and September, there were reports of mass mortality of fauna, from Cancun to Xcalak (Supp. Fig. 2), in association with massive arrival of *Sargassum* spp. and the subsequent formation of *Sargassum*-brown-tides. It should be noted, however, that these reports came mostly from tourist zones, where beach clean-up is frequent, so it is likely that the number of dead species is an underestimate. These reports were incorporated into a database by Universidad Nacional Autónoma de México, El Colegio de la Frontera Sur and the Puerto Morelos Reef National Park. These data show that individuals belonging to 78 faunal species were recorded in association with the *Sargassum* event of 2018. Fish were most commonly affected (59%) followed by Crustacea (28%), Echinoderms (5%), Mollusca (4%), and Polychaeta (4%) (Fig. 1). The majority of the reports ( $N = 75$ ) were for the northern sector of the Mexican Caribbean coast, compared to the center ( $N = 9$ ) and south ( $N = 8$ ), where there is less tourism (and therefore less beach cleaning).

Of the 46 species of fish that were recorded, 80% have a demersal neritic habitat and are commonly found on coral reefs (cf. Claro and Robertson, 2010), while all dead organisms belonging to other faunal groups were benthic (Table 1). Only four of the recorded fish species (*Haemulon* sp., *Harengula* sp., *Caranx* sp. and *Abudefduf saxatilis*) were reported for all three sectors of the coast (Table 1). Other fauna that were frequently reported dead were the arenicolous sea cucumber *Holothuria arenicola*, the fish *Diodon holacanthus*, morays of the genera *Gymnothorax*, the crabs *Arenaeus cribarius* and *Callinectes similis* and the mollusk *Octopus* sp.

In most cases, dead organisms beached individually or in small (< 10) quantities, but on five occasions mass fish mortality (> 100 individuals) was reported at Punta Caracol, Puerto Morelos (May 2018; Fig. 2), Mahahual (July and August 2018) and Xcalak (August and

September 2018). In these reports, dead organisms were mainly sardines of the family Clupeidae genera *Harengula*, in the northern and southern sectors, and grunts of the genera *Haemulon*, in the southern sector (Table 1).

The cause of faunal mortality recorded in the 2018 *Sargassum* event appears to be hypoxia and deterioration of water quality. All mass mortality events were recorded on beaches adjacent to shallow (< 3 m) reef lagoons, during calm windless days, and after several days of *Sargassum* spp. build-up on the shore. Analysis of water samples ( $N = 7$ , Supp. Table 2), taken between 45 and 480 m from the shore at one of these sites (Punta Caracol, Puerto Morelos), two days after the massive mortality of dead fauna (May 9th, 2018, Fig. 2), showed low concentration of dissolved oxygen (Mean:  $2.9 \text{ mg L}^{-1}$ ; range:  $1.9\text{--}4.2 \text{ mg L}^{-1}$ ) and high values of ammonium (mean:  $6.0 \text{ } \mu\text{mol L}^{-1}$ ; range:  $4.6\text{--}8.8 \text{ } \mu\text{mol L}^{-1}$ ) and phosphorus (mean:  $5.0 \text{ } \mu\text{mol L}^{-1}$ ; range:  $3.9\text{--}6.12 \text{ } \mu\text{mol L}^{-1}$ ) as far as 480 m from the shore (Supp. Table 1). Oxygen values were lower than those recorded in Puerto Morelos reef lagoon in 2014, before the massive arrival of *Sargassum* spp. (mean:  $5.35$ , SD:  $0.22 \text{ mg L}^{-1}$ ), and ammonium and phosphorus values were an order of magnitude higher than the typical values for this area ( $0.03\text{--}0.24 \text{ } \mu\text{mol L}^{-1}$  and  $0.32\text{--}0.57 \text{ } \mu\text{mol L}^{-1}$  respectively; Almazán-Becerril et al., 2014). Marine organisms are sensitive to low levels of dissolved oxygen; their metabolism is affected at levels below  $4.0 \text{ mg O}_2 \text{ L}^{-1}$ , and mortality ensues in fish and crustaceans when it drops below  $2.0 \text{ mg O}_2 \text{ L}^{-1}$  (Vaquer-Sunyer and Duarte, 2008). Also, high ammonium levels are toxic in aquatic environments (Gray et al., 2002). Although sulfide was not measured in these samples, we hypothesize that the high temperature ( $28.2\text{--}29.6 \text{ } ^\circ\text{C}$ ) and the large influx of organic matter from decomposing *Sargassum* spp. led to rapid reduction in dissolved oxygen and high levels of  $\text{H}_2\text{S}$ . Sulfide is toxic for most aerobic organisms, even at low concentrations (Janas and Szaniawska, 1996), as it diffuses through respiratory membranes and inhibits cytochrome oxidase function (cytochrome *c* oxidase; cytochrome *a*, *a*3), even in short-term exposures, especially when water temperature is high (Torrans and Clemens, 1982). In fish, sublethal exposure to sulfide has been found to induce gill and liver damage (Kiemer et al., 1995) and even in tolerant organisms (i.e. polychaetes) sulfide induces damage to RNA and DNA in coelomocytes and epithelial tissue (Joyner-Matos et al., 2010).

In conclusion, the mortality of fauna associated with the 2018 *Sargassum* event was widespread along the Mexican Caribbean coast and affected individuals of a large number of species, mostly of fish with demersal neritic habitats and crustaceans. In the absence of other algal blooms or major disturbances (i.e. storms) it appears that the combined effect of high ammonium and  $\text{H}_2\text{S}$  concentrations together with hypoxic conditions were responsible for this mortality event. It is uncertain how much of the lagoon water column was affected, but if massive influx of *Sargassum* spp. becomes an annual event it could potentially have a deleterious impact on the already degraded coral reefs of the region. This could be further exacerbated by the lagoonal placement of interception barriers to prevent the *Sargassum* spp. from reaching shore if it is left to sink and die in the lagoon. Different technologies for the removal of *Sargassum* spp. from the barriers are being developed, including the use of “Sargaboats” and pumps. Another option is to use barriers to deflect the algae to locations on the beach where collection and transport is easier. Management of beaches adjacent to coral reef lagoons that are receiving massive amounts of *Sargassum* spp. should consider these potential problems before deploying such barriers.

**Table 1**

Fauna species observed dead on beaches of the Mexican Caribbean coast during the massive influx of pelagic *Sargassum* spp. of 2018. The habitat of each species is indicated. Sectors: North sector: From Cancun to Puerto Aventuras, Central sector: Sian Ka'an, South sector: From Mahahual to Xcalak. Habitat determined after [Claro et al. \(2014\)](#) for fish and [Felder and Camp \(2009\)](#) for Crustacean. M: organisms observed dead in massive amounts (> 100).

Group	Species	Habitat	Sector		
			North	Central	South
Fish	<i>Abudefduf saxatilis</i>	Demersal Neritic	*	*	*
	<i>Acanthostracion quadricornis</i>	Demersal Neritic	*		
	<i>Acanthostracion polygonius</i>	Demersal Neritic	*		
	<i>Callechelys bilinearis</i>	Demersal Neritic	*		
	<i>Canthigaster rostrata</i>	Demersal Neritic	*	*	
	<i>Canthigaster</i> sp.	Demersal Neritic	*		
	<i>Caranx</i> sp.	Coastal Epipelagic	*	*	*
	<i>Diodon holocanthus</i>	Demersal Neritic	*		*
	<i>Enchelycore carychroa</i>	Demersal Neritic	*		
	<i>Enchelycore nigricans</i>	Demersal Neritic	*		
	<i>Gymnothorax funebris</i>	Demersal Neritic	*		
	<i>Gymnothorax moringa</i>	Demersal Neritic	*		
	<i>Gymnothorax</i> sp.	Demersal Neritic	*	*	
	<i>Haemulon aurolineatum</i>	Demersal Neritic	*		
	<i>Haemulon flavolineatum</i>	Demersal Neritic	*		
	<i>Haemulon parra</i>	Demersal Neritic	*		
	<i>Haemulon sciurus</i>	Demersal Neritic	*		
	<i>Haemulon</i> sp.	Demersal Neritic	*	*	M
	<i>Halichoeres bivittatus</i>	Demersal Neritic		*	
	<i>Harengula jaguana</i>	Demersal Pelagic			*
	<i>Harengula</i> sp.	Coastal Pelagic	M	*	M
	<i>Hemiramphus balao</i>	Coastal Epipelagic	*		
	<i>Histrio histrio</i>	Epipelagic	*		
	<i>Kyphosus sectatrix</i>	Pelagic Neritic	*		
	<i>Lactophrys bicaudalis</i>	Demersal Neritic	*		
	<i>Lactophrys trigonus</i>	Demersal Neritic	*		
	<i>Lutjanus analis</i>	Demersal Neritic	*		
	<i>Lutjanus apodus</i>	Demersal Neritic	*		
	<i>Megalops atlanticus</i>	Pelagic Neritic	*		
	<i>Mycteroperca bonaci</i>	Demersal Neritic			*
	<i>Microspathodon chrysurus</i>	Demersal Neritic	*		
	<i>Ocyurus chrysurus</i>	Demersal Neritic	*		
	<i>Paraclinus</i> sp.	Demersal Neritic	*		
	<i>Polydactylus virginicus</i>	Demersal Neritic	*		
	<i>Pomacanthus arcuatus</i>	Demersal Neritic	*		
	<i>Scorpaena</i> sp.	Demersal Neritic	*	*	
	<i>Selene vomer</i>	Demersal Neritic	*		
	<i>Sparisoma aurofrenatum</i>	Demersal Neritic	*		
	<i>Sparisoma chrysopterygum</i>	Demersal Neritic	*		
	<i>Sparisoma rubripinne</i>	Demersal Neritic	*		
	<i>Sparisoma radians</i>	Demersal Neritic	*		
	<i>Sparisoma</i> sp.	Demersal Neritic	*	*	
	<i>Sphoeroides testudineus</i>	Demersal Neritic	*		
	<i>Styracura schmardae</i>	Demersal Neritic	*		
	<i>Trachinotus falcatus</i> (juvenile)	Pelagic Epibenthic	*		*
	<i>Urobatis jamaicensis</i>	Demersal	*		
	Crustacean	<i>Achelous spinimanus</i>	Benthic	*	
<i>Alpheus heterochaelis</i>		Benthic	*		
<i>Ampithoe</i> sp.		Benthic	*		
<i>Arenaeus cribarius</i>		Benthic	*		
<i>Callinectes marginatus</i>		Benthic	*		
<i>Callinectes similis</i>		Benthic	*		
<i>Carpas algicola</i>		Benthic	*		
<i>Cataleptodius floridanus</i>		Benthic	*		
<i>Chorinus heros</i>		Benthic	*		
<i>Cronius ruber</i>		Benthic	*		
<i>Gonodactylus</i> sp.		Benthic	*		
<i>Leander tenuicornis</i>		Benthic	*		
<i>Lysmata</i> sp.		Benthic	*		
<i>Menippe nodifrons</i>		Benthic	*		
<i>Mithrax forceps</i>		Benthic	*		
<i>Panulirus argus</i>		Benthic	*		
Penaeidae sp.		Benthic	*		
<i>Portunus</i> sp.		Benthic	*		
<i>Portunus sayi</i>		Benthic	*		
<i>Pseudosquilla</i> sp.		Benthic	*		
<i>Stenopus hispidus</i>		Benthic	*		
<i>Stenorhynchus seticornis</i>		Benthic	*		

(continued on next page)

Table 1 (continued)

Group	Species	Habitat	Sector		
			North	Central	South
Echinoderms	<i>Echinometra</i> sp.	Benthic	*		
	<i>Holothuria arenicola</i>	Benthic	*		
	<i>Ophioderma</i> sp.	Benthic	*		
	<i>Meoma ventricosa</i>	Benthic	*		
Molluscs	<i>Cyphoma gibbosum</i>	Benthic	*		
	<i>Litiopa melanostoma</i>	Benthic	*		
	<i>Octopus</i> sp.	Benthic	*		
Polychaetes	<i>Eupolymnia crassicornis</i>	Benthic	*		
	<i>Hermodice carunculata</i>	Benthic	*		
	Familia Syllidae	Benthic	*		



Fig. 2. Mass mortality of fauna associated with Sargassum-brown-tide at Punta Caracol, Puerto Morelos, in May 2018. a. Onshore accumulation of *Sargassum* spp. and *Sargassum*-brown-tide, b. Dead fish scattered along the beach, c. Different fish species found dead on the beach, d. Fish, mostly of the genera *Harengula*, that died in massive amounts. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

## CRedit authorship contribution statement

**R.E. Rodríguez-Martínez:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Visualization, Writing - original draft, Writing - review & editing. **A.E. Medina-Valmaseda:** Data curation, Investigation, Writing - original draft, Writing - review & editing. **P. Blanchon:** Writing - original draft, Writing - review & editing. **L.V. Monroy-Velázquez:** Data curation, Investigation, Writing - original draft, Writing - review & editing. **A. Almazán-Becerril:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing - original draft, Writing - review & editing. **B. Delgado-Pech:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing - original draft, Writing - review & editing. **L. Vázquez-Yeomans:** Conceptualization, Data curation, Investigation, Writing - original draft, Writing - review & editing. **V. Francisco:** Conceptualization, Data curation, Investigation, Visualization, Writing - original draft, Writing - review & editing. **M.C. García-Rivas:** Conceptualization, Writing - original draft, Writing - review & editing.

## Acknowledgments

Special thanks to Erika Hernández, Horacio Ocampo, Verónica Ramos, Alfredo Arellano, Luis Ramos, Enrique Gallegos, Ma. Jesús Hermosillo, Guadalupe Paz, Yazmin Novelo, Elva Leyva, Fortunato Herrera, Cedrella Medina, Roberto Herrera, Susana Rivero, Jully Novelo and Jorge Peniche for sharing the photographs of dead fauna associated to *Sargassum*-brown-tides and to Antonio Ortiz, Gerardo Castañeda, Antonio Lascano, Imelda Juárez, Lucia Paredes, Miguel A. Diego and Verónica Ramos for sharing the data of the volume of *Sargassum* spp. removed by hotels from the beach in 2018. The authors claim no conflict of interest with this work.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.marpolbul.2019.06.015>.

## References

- Adet, L., Nsofor, G.N., Ogunjobi, K.O., Camara, B., 2017. Knowledge of climate change and the perception of Nigeria's coastal communities on the occurrence of *Sargassum natans* and *Sargassum fluitans*. Open Access Library Journal 4 <https://doi.org/10.4236/oalib.1104198>. e4198.
- Almazán-Becerril, A., Abundes, M., Francisco, V., López, I., Ramos, L., Álvarez-Filip, L., Delgado-Pech, A., Escobar-Morales, S., Gómez-Campo, K., López-Lodoño, T., Irola-Sansores, E.D.B., Manzo, S., 2014. Programa de Monitoreo Biológico (PROMOBI) – 2014. Informe Final. CONANP (88 p).
- Anderson, D.M., 2007. The ecology and oceanography of harmful algal blooms: multi-disciplinary approaches to research and management. In: IOC Technical Series. vol. 74 UNESCO (IOC/2007/TS/74).
- Claro, R., Robertson, D.R., 2010. Los peces de Cuba. Instituto de Oceanografía. CITMA, La Habana, Cuba 978-959-298-019-8 en CD-ROM.
- Claro, R., Lindeman, K.C., Parenti, L.R. (Eds.), 2014. Ecology of the Marine Fishes of Cuba. Smithsonian Institution Press, Washington and London (253 p).
- Felder, D.L., Camp, D.K. (Eds.), 2009. Gulf of Mexico Origin, Waters, and Biota. vol. 1 Texas A&M University Press, College Station, Tex (Biodiversity. 1384 p).
- Gower, J., Young, E., King, S., 2013. Satellite images suggest a new *Sargassum* source region in 2011. Remote Sensing Letters 4 (8), 764–773. <https://doi.org/10.1080/2150704X.2013.796433>.
- Gray, J.S., Shiu-sun, R., Or, Y.Y., 2002. Effects of hypoxia and organic enrichment on the coastal marine environment. Mar. Ecol. Prog. Ser. 238, 249–279. <https://doi.org/10.3354/meps238249>.
- Harvell, C.D., Kim, K., Burkholder, J.M., Colwell, R.R., Epstein, P.R., Grimes, D.J., Hofmann, E., Lipp, E.K., Osterhaus, A.D.M.E., Overstreet, R.M., Porter, J.W., Smith, G.W., Vasta, G.R., 1999. Emerging marine diseases—climate links and anthropogenic factors. Science 285 (5433), 1505–1510. <https://doi.org/10.1126/science.285.5433.1505>.
- Hsieh, H.L., Hsien, Y.-L., Jeng, M.S., Tsai, W.S., Su, W.C., Chen, C.A. 2008. Tropical fishes killed by the cold. Coral Reefs, 27: 599. <https://doi.org/10.1007/s00338-008-0378-3>.
- Janas, U., Szaniawska, A., 1996. The influence of hydrogen sulphide on macrofaunal biodiversity in the Gulf of Gdansk. Oceanologia 38, 127–142 (PL ISSN 0078–3234).
- Johnson, D.R., Ko, D.S., Franks, J.S., Moreno, P., Sánchez-Rubio, G., 2013. The Sargassum invasion of the Eastern Caribbean and dynamics of the equatorial North Atlantic. In: Proceed. 65th GCFI, Santa Marta, Colombia, pp. 102–103.
- Joyner-Matos, J., Predmore, B.L., Stein, J.R., Leeuwenburgh, C., Julian, D., 2010. Hydrogen sulfide induces oxidative damage to RNA and DNA in a sulfide-tolerant marine invertebrate. Physiol. Biochem. Zool. 83, 356–365. <https://doi.org/10.1086/597529>.
- Justić, D., Rabalais, N.N., Turner, R.E., 1996. Effects of climate change on hypoxia in coastal waters: a doubled CO<sub>2</sub> scenario for the northern Gulf of Mexico. Limnol. Oceanogr. 41 (5), 992–1003. <https://doi.org/10.4319/lo.1996.41.5.0992>.
- Kierner, M.C.B., Black K.D., Lussot D., Bullock A.I., Ezzi I. 1995. The effects of chronic and acute exposure to hydrogen sulphide on Atlantic salmon (*Salmo salar* L.). Aquaculture, 135: 311–327. [https://doi.org/10.1016/0044-8486\(95\)01025-4](https://doi.org/10.1016/0044-8486(95)01025-4).
- Laffoley, D.d'A., Roe, H.S.J., Angel, M.V., Ardron, J., Bates, N.R., Boyd, I.L., Brooke, S., Buck, K.N., Carlson, C.A., Causey, B., Conte, M.H., Christiansen, S., Cleary, J., Donnelly, J., Earle, S.A., Edwards, R., Gjerde, K.M., Giovannoni, S.J., Gulick, S., Gollock, M., Hallett, J., Halpin, P., Hanel, R., Hemphill, A., Johnson, R.J., Knap, A.H., Lomas, M.W., McKenna, S.A., Miller, M.J., Miller, P.I., Ming, F.W., Moffitt, R., Nelson, N.B., Parson, L., Peters, A.J., Pitt, J., Rouja, P., Roberts, J., Roberts, J., Seigel, D.A., Siuda, A.N.S., Steinberg, D.K., Stevenson, A., Sumaila, V.R., Swartz, W., Thorrold, S., Trott, T.M., Vats, V., 2011. The protection and management of the Sargasso Sea: the golden floating rainforest of the Atlantic Ocean. In: Summary Science and Supporting Evidence Case. Sargasso Sea Alliance (44 p).
- Lapointe, B.E., 1997. Nutrient thresholds for bottom-up control of macroalgal blooms on coral reefs in Jamaica and southeast Florida. Limnol. Oceanogr. 42, 1119–1131. [https://doi.org/10.4319/lo.1997.42.5\\_part\\_2.1119](https://doi.org/10.4319/lo.1997.42.5_part_2.1119).
- Lessios, H.A., Cubit, J.D., Robertson, D.R., Shulman, M.J., Parker, M.R., Garrity, S.D., Levings, S.C., 1984. Mass mortality of *Diadema antillarum* on the Caribbean coast of Panama. Coral Reefs 3, 173–182. <https://doi.org/10.1007/BF00288252>.
- Oyesiku, O.O., Egunyomi, A., 2014. Identification and chemical studies of pelagic masses of *Sargassum natans* (Linnaeus) Gaillon and *S. fluitans* (Borgesen) Borgesen (brown algae), found offshore in Ondo state, Nigeria. Afr. J. Biotechnol. 13, 1188–1193. <https://doi.org/10.5897/AJB2013.12335>.
- Pinheiro, H.T., Gasparini, J.L., Joyeux, J.C., 2010. Reef fish mass mortality event in an isolated island off Brazil, with notes on recent similar events at Ascension, St. Helena and Maldives. Marine Biodiversity Records 3, 1–4. <https://doi.org/10.1017/S1755267210000424>.
- Rodríguez-Martínez, R.E., van Tussenbroek, B.I., Jordán-Dahlgren, E., 2016. Afluencia masiva de sargazo pelágico a la costa del Caribe mexicano (2014–2015). In: García-Mendoza, E., Quijano-Scheggia, S.I., Olivares-Ortiz, A., Núñez-Vázquez, E.J. (Eds.), Florecimientos Algales nocivos en México. CICESE, Ensenada, México (ISBN: 352–365.978-607-95688-5-6).
- Sankaré, Y., Komoé, K., Aka, K.S., Fofié, N.B.Y., Bamba, A., 2016. Répartition et abondance des sargasses *Sargassum natans* et *Sargassum fluitans* (Sargassaceae, Fucales) dans les eaux marines ivoiriennes (Afrique de l'Ouest). International Journal of Biological and Chemical Sciences 10, 1853–1856. <https://doi.org/10.4314/ijbcs.v10i4.33>.
- Sissini, M.N., Barbosa de Barros Barreto, M.B., Menezes Széchi, M.T., Bouças de Lucena, M., Cabral Oliveira, M., Gower, J., ... Antunes Horta, P., 2017. The floating *Sargassum* (Phaeophyceae) of the South Atlantic Ocean—likely scenarios. Phycologia 56, 321–328. <https://doi.org/10.2216/16-92.1>.
- Smetacek, V., Zingone, A., 2013. Green and golden seaweed tides on the rise. Nature 504, 84–88.
- Torrans, E.L., Clemens, H.P., 1982. Physiological and biochemical effects of acute exposure of fish to hydrogen sulfide. Comparative Biochemistry and Physiology Part C: Comparative Pharmacology 71 (2), 183–190. [https://doi.org/10.1016/0306-4492\(82\)90034-X](https://doi.org/10.1016/0306-4492(82)90034-X).
- van Tussenbroek, B.I., Hernández Arana, H.A., Rodríguez-Martínez, R.E., Espinoza-Ávalos, J., Canizales-Flores, H.M., González-Godoy, C.E., Barba-Santos, M.G., Vega-Zepeda, A., Collado-Vides, L., 2017. Severe impacts of brown tides caused by *Sargassum* spp. on near-shore Caribbean seagrass communities. Mar. Pollut. Bull. 122, 272–278. <https://doi.org/10.1016/j.marpolbul.2017.06.057>.
- Vaquer-Sunyer, R., Duarte, C., 2008. Thresholds of hypoxia for marine biodiversity. Proc. Natl. Acad. Sci. U. S. A. 105 (40), 15452–15457. <https://doi.org/10.1073/pnas.0803833105>.